EXHIBIT 4



Can dehydration cause creatinine to rise? If so, by what factor?

Post date: June 24, 2013

Dehydration does cause the serum creatinine to rise and the estimated glomerular filtration rate (eGFR) will, accordingly, fall. The degree of change is generally proportional to the degree of dehydration. Severe dehydration can actually cause acute kidney injury and may lead to a need for dialysis therapy.

- My blood creatinine of 1.60 mg/dL, age 72, male, diabetic with HbA1c of 5.5%. Does this indicate CKD and if so, what stage and what should I do? I live in Indonesia but am American.
- I had a UTI infection in April. I did not get better. I was ill had burning through out my body. Then treated for a kidney infection. They are calling my burning chronic pain? I am still being treated for kidney infection and then bladder 6 treatments since May? I have lost 22 pounds since May, still have burning. Strange bowel movements, swelling in groin, inner thigh and buttock?
- Are there any solutions for continued calf, ankle and foot swelling? My numbers are back to normal and I try to monitor salt intake, but it's been a year and they look awful. I also have very strong muscle cramps, sometimes in the groin area and some times the calf. I take magnesium and Co Q10 every day. I came off Prednisone 2 months ago and still get very achy joints and stiff muscles. Takes a few moments to get started after sitting for awhwile. My heart is good and I'm anxious to get back to working out without additional cramping! Have lost 11 lbs with 40 more to go.

Privacy policy
Legal & copyright
State fundraising notices

30 East 33rd Street, New York, NY 10016 © 2023 National Kidney Foundation Inc.



C Log in





Patient Care & Health Information

Tests & Procedures

Blood urea nitrogen (BUN) test

Request an Appointment

About

Print

Advertisement

Overview

A common blood test, the blood urea nitrogen (BUN) test reveals important information about how well your kidneys are working. A <u>BUN</u> test measures the amount of urea nitrogen that's in your blood.

Here's how your body typically forms and gets rid of urea nitrogen:

- Your liver produces ammonia which contains nitrogen after it breaks down proteins used by your body's cells.
- The nitrogen combines with other elements, such as carbon, hydrogen and oxygen, to form urea, which is a chemical waste product.
- The urea travels from your liver to your kidneys through your bloodstream.
- Healthy kidneys filter urea and remove other waste products from your blood.
- The filtered waste products leave your body through urine.

A <u>BUN</u> test can reveal whether your urea nitrogen levels are higher than normal, suggesting that your kidneys may not be working properly.

Products & Services

Book: Mayo Clinic Family Health Book, 5th Edition

Show more products from Mayo Clinic

Mayo Clinic does not endorse companies or products. Advertising revenue supports our $n\alpha$ for-profit mission.

Advertising & Sponsorship

Policy | Opportunities | Ad Choices

Mayo Clinic Press

Check out these best-sellers and special offers on books and newsletters from May Clinic Press.

Mayo Clinic on Incontinence

NEW - Future Care

Mayo Clinic on Hearing and Balance

FREE Mayo Clinic Diet Assessment

Mayo Clinic Health Letter - FREE book

You may need a BUN test:

- If your doctor suspects that you have kidney disease or damage
- If your kidney function needs to be evaluated, especially if you have a chronic condition such as diabetes or high blood pressure
- To help determine the effectiveness of dialysis treatment if you're receiving hemodialysis or peritoneal dialysis
- As part of a blood test group to help diagnose a number of other conditions, such as liver damage, urinary tract obstruction, congestive heart failure or gastrointestinal bleeding — although an abnormal <u>BUN</u> test result alone doesn't confirm any of these conditions

If kidney problems are the main concern, the creatinine levels in your blood will likely also be measured when your blood is tested for urea nitrogen levels. Creatinine is another waste product that healthy kidneys filter out of your body through urine. High levels of creatinine in your blood may be a sign of kidney damage.

Your doctor may also test how well your kidneys are removing waste from the blood. To do this, you may have a blood sample taken to calculate your estimated glomerular filtration rate (GFR). The <u>GFR</u> estimates the percentage of kidney function you have.

More Information

Diabetic coma

Request an appointment

From Mayo Clinic to your inbox



Sign up for free, and stay up to date of presearch advancements, health tips and current health topics, like COVID-19, plus expertise on managing health. Click here for an email preview.

Enter your email

Learn more about Mayo Clinic's use of data.

How you prepare

If your blood sample is being tested only for <u>BUN</u>, you can eat and drink normally before the test. If your blood sample will be used for additional tests, you may need to fast for a certain amount of time before the test. Your doctor will give you specific instructions.

What you can expect

During the <u>BUN</u> test, a member of your health care team takes a sample of blood by inserting a needle into a vein in your arm. The blood sample is sent to a lab for analysis. You can return to your usual activities immediately.

Results

Results of the <u>BUN</u> test are measured in milligrams per deciliter (mg/dL) in the United States and in millimoles per liter (mmol/L) internationally. In general, around 6 to 24 mg/dL (2.1 to 8.5 mmol/L) is considered normal.

But normal ranges may vary, depending on the reference range used by the lab and your age. Ask your doctor to explain your results.

Urea nitrogen levels tend to increase with age. Infants have lower levels than other people do, and the range in children varies.

Generally, a high <u>BUN</u> level means your kidneys aren't working well. But elevated <u>BUN</u> can also be due to:



- Dehydration, residency fram (กษณ์ และ คระบาร์ คร
- · Urinary tract obstruction
- · Congestive heart failure or recent heart attack
- Gastrointestinal bleeding
- Shock
- Severe burns
- · Certain medications, such as some antibiotics
- A high-protein diet

If kidney damage is a concern, ask your doctor what factors may be contributing to the damage and what steps you can take to try to control them.

By Mayo Clinic Staff

Request an appointment

Share on: <u>Facebook</u> <u>Twitter</u> <u>Print</u> Aug. 19, 2021

Show references ∨

Related

Diabetic coma

Products & Services

Book: Mayo Clinic Family Health Book, 5th Edition

Show more products and services from Mayo Clinic

Blood urea nitrogen (BUN) test



Your gift holds great power – donate today!

Make your tax-deductible gift and be a part of the cutting-edge research and care that's changing medicine.

	Donate	
Find a doctor		
Explore careers		
Sign up for free e-newsletters		

About Mayo Clinic

About this Site

Contact Us

Locations

Health Information Policy

Media Requests

News Network

Medical Professionals

AskMayoExpert

Clinical Trials

Mayo Clinic Alumni Association

Refer a Patient



Case 5:22-cv-00752-F Document 29-4 Filed 06/19/23 Page 8 of 9

Businesses

Executive Health Program

International Business Collaborations

Facilities & Real Estate

Supplier Information

Students

Admissions Requirements

Degree Programs

Student & Faculty Portal

Researchers

Research Faculty

Laboratories

International Patients

Appointments

Financial Services

International Locations & Offices

Charitable Care & Financial Assistance

Community Health Needs Assessment

Financial Assistance Documents - Arizona

Financial Assistance Documents - Florida

Financial Assistance Documents - Minnesota

Follow Mayo Clinic



Terms and Conditions

Privacy Policy

Notice of Privacy Practices

Notice of Nondiscrimination

Advertising and Sponsorship Policy

Site Map

Language: English

© 1998-2023 Mayo Foundation for Medical Education and Research (MFMER). All rights reserved.

